

## **Seattle Genova**

Tel: +1 (425) 247-3088 Fax: +1 (425) 650-9990

Email: info@seattle-genova.com Web: www.seattle-genova.com Address: 18110 SE 34TH ST STE 455, Vancouver, WA 98683

## **Rasburicase In Vitro Transcribed mRNA-LNP**

Catalog Number:SG-MRNA-LNP-1922

DESCRIPTION	
Product Name	Rasburicase In Vitro Transcribed mRNA-LNP
Gene Name	Rasburicase
Source	The ORF of Rasburicase was cloned in our IVT vector and mRNA was prepared through in vitro transcription and purification. The purified mRNA was further encapsulated with LNP(DSPC:Cholesterol:DMG-PEG:SM102).
Alternative names	Rasburicase
SPECIFICATIONS	
Cap	m7GpppN
5'-UTR	5' -untranslated region derived from human alpha-globin RNA with an optimized Kozak sequence
ORF	Rasburicase
3'-UTR	3' UTR comprising two sequence elements derived from the aminoterminal enhancer of split (AES) mRNA and the mitochondrial encoded 12S ribosomal RNA
Poly(A) Tail	A 110-nucleotide poly(A)-tail consisting of a stretch of 30 adenosine residues, followed by a 10-nucleotide linker sequence and another 70 adenosine residues.
Modifications	N1-methyl-pseudouridine
Neutral Lipid	1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC)
Cholesterol	Cholesterol
Lonizable Lipid	1,2-dimyristoyl-rac-glycero-3-methoxypolyethylene glycol-2000 (PEG2000-DMG)
PEG-lipid	Heptadecan-9-yl 8-((2-hydroxyethyl)(8-(nonyloxy)—8-oxooctyl)amino)octanoate)(SM-102)
Storage	-20 °C
Buffer	PBS, pH7.4
Cryoprotectant	Trehalose
BACKGROUND	
Gene Accession	
Gene Alias	Rasburicase
Background	Rasburicase catalyzes enzymatic oxidation of uric acid into an inactive and soluble metabolite (allantoin).